arXiv:1607.02590v1 [math.RA] 9 Jul 2016

# APPLICATIONS OF THE WALL FORM TO UNIPOTENT ISOMETRIES OF INDEX TWO

A.-H. NOKHODKAR

Abstract. We investigate the Wall form of unipotent elements of index two in the orthogonal group and obtain a decomposition for these elements. Also, in characteristic two, the relation between the Wall form and some invariants of the induced involution on the Clifford algebra is studied.
**Keywords.** Wall form, unipotent isometry, algebra with involution, Clifford algebra.
**Mathematics Subject Classification.** 11E88, 11E39, 16W10.

## 1. Introduction

The *Wall form* of an isometry is a nondegenerate bilinear form defined on its residual space (see [17, pp. 12–13]). According to [16, Theorem 1.3.1], isometries of a quadratic space can be classified, up to conjugation, by their Wall forms. The Wall form was applied in [5] and [7] in connection with the Zassenhaus decomposition [19] of isometries in characteristic $\neq 2$. It was also used in [6] to study the length of elements in the commutator subgroup of the orthogonal group. In characteristic 2, this form was applied in [1] to study the relation between the commutator subgroup of the orthogonal group and the spinorial kernel of a nondegenerate defective quadratic space.

Let $(V, q)$ be a quadratic space over a field $F$. An isometry $\tau$ of $(V, q)$ is called *unipotent* if $(\tau - \mathrm{id})^k = 0$ for some nonnegative integer $k$. The minimum integer $k$ with this property is called the *unipotency index* of $\tau$. According to [4, (6.3)], an isometry $\tau$ has unipotency index 2 if and only if its residual space is totally singular. Also, unipotent isometries of index 2 are exactly those isometries whose Wall forms are antisymmetric (see [5, p. 116]).

In this work we investigate some applications of the Wall form to unipotent elements of index 2 in the orthogonal group. We start with some observations on *indecomposable* isometries (i.e., isometries which admit no invariant regular proper subspace). In characteristic two, every unipotent isometry of index 2 is an involution in the orthogonal group. Such an involution is indecomposable if and only if it is either a 2-dimensional reflection or a 4-dimensional *basic null involution* (see [18, Theorem 1]). Some properties of 4-dimensional basic null involutions, called *interchange isometries* are investigated in [13].

In §3, we generalize the definition of interchange isometries to arbitrary characteristic. As we shall see in (3.5), a 4-dimensional isometry is an inetrchange isometry if and only if it is indecomposable and unipotent of index 2. Also, using the Wall form we shall obtain in (3.10) a decomposition of unipotent isometries of index 2 into orthogonal sums of interchange isometries (and 2-dimensional reflections if $\operatorname{char} F = 2$). In characteristic two, using the Jordan canonical form of a matrix, this decomposition was already obtained in [18, Theorem 2]. However, the connection between this decomposition and the Wall form has not been studied.

Let $(V, q)$ be a quadratic space over a field $F$. Every involution $\tau$ in $O(V, q)$ induces a *natural* involution $J_\tau$, introduced by D. B. Shapiro [14], on the Clifford

algebra $C(q)$ satisfying $J_\tau(v) = \tau(v)$ for $v \in V$. In [11], it was shown that every totally decomposable algebra with involution over a field of characteristic different from 2 can be expressed as the Clifford algebra of a quadratic space with a natural involution induced by an involution in the orthogonal group (see [13] for a characteristic 2 counterpart). In characteristic 2, it is readily seen that the converse is also true (see (4.5)). Thus, if $\operatorname{char} F = 2$, there exists a natural correspondence between totally decomposable algebras with involution and Clifford algebras with involution induced by an involution in the orthogonal group.

In [2] and [12] some invariants of a totally decomposable algebra with orthogonal involution $(A, \sigma)$ in characteristic 2, such as the *Pfister invariant* and the subalgebra $\Phi(A, \sigma)$ were introduced. In [12, (6.5)] it was shown that totally decomposable orthogonal involutions on a given algebra can be classified by their Pfister invariants. It was also shown that the Pfister invariant of $(A, \sigma)$ is isometric to an associative bilinear form defined naturally on $\Phi(A, \sigma)$ (see [12, (5.5)]). In view of these characterizing properties, it would be interesting to study the relation between these invariants and the Wall form of $\tau$, when $(A, \sigma)$ is expressed as a Clifford algebra with natural involution $J_\tau$. This investigation, which is the main object of §4, gives a geometric interpretation of the aforementioned invariants. As a final application, we obtain in (4.8), necessary and sufficient conditions for $(C(q), J_\tau)$ to be isomorphic to the matrix algebra with the transpose involution $(M_{2^n}(F), t)$.

## 2. Preliminaries

Let $V$ be a finite dimensional vector space over a field $F$. A bilinear form $\mathfrak{b}$ on $V$ is called *alternating* if $\mathfrak{b}(u, u) = 0$ for every $u \in V$. Otherwise, $\mathfrak{b}$ is called *nonalternating*. Every nonalternating symmetric bilinear space $(V, \mathfrak{b})$ has an *orthogonal basis*, i.e., a basis $(v_1, \cdots, v_n)$ such that $\mathfrak{b}(v_i, v_j) = 0$ for every $i \neq j$ (see [3, (1.17)]). In this case, $\mathfrak{b}$ is denoted by $\langle \alpha_1, \cdots, \alpha_n \rangle$, where $\alpha_i = \mathfrak{b}(v_i, v_i) \in F$. The form $\langle\langle \alpha_1, \cdots, \alpha_n \rangle\rangle := \langle 1, \alpha_1 \rangle \otimes \cdots \otimes \langle 1, \alpha_n \rangle$ is called a *bilinear $n$-fold Pfister form*. For a bilinear form $\mathfrak{b}$ we denote the form $\mathfrak{b} \perp \cdots \perp \mathfrak{b}$ ($r$ times) by $r\mathfrak{b}$.

Let $q : V \to F$ be a quadratic form. The bilinear form $\mathfrak{b}_q : V \times V \to F$ defined by $\mathfrak{b}_q(u, v) = q(u + v) - q(u) - q(v)$ is called the *polar form* of $q$. A subspace $W$ of $V$ is called *regular* if $W \cap W^\perp = \{0\}$, where $W^\perp = \{x \in V \mid b_q(x, y) = 0 \text{ for all } y \in W\}$. Otherwise, $W$ is called *singular*. The pair $(V, q)$ is called a *quadratic space* if $V$ is regular. A quadratic form $q$ is called *totally singular* if $\mathfrak{b}_q = 0$. Nontrivial totally singular quadratic forms only exist in characteristic two. A nonzero vector $v \in V$ is called *isotropic* if $q(v) = 0$. A subspace $W$ of $V$ is called *totally isotropic* if all nonzero vectors of $W$ are isotropic.

Let $V$ be a $2n$-dimensional vector space over a field $F$ and let $\lambda = \pm 1$. A bilinear form $\mathfrak{b}$ (resp. a quadratic form $q$) on $V$ is called *hyperbolic* if it has a *hyperbolic basis*, i.e., a basis $\mathcal{B} = (u_1, v_1, \cdots, u_n, v_n)$ such that: (i) $\mathfrak{b}(x, x) = 0$ (resp. $q(x) = 0$) for every $x \in \mathcal{B}$; (ii) $\mathfrak{b}(u_i, v_i) = \lambda\mathfrak{b}(v_i, u_i) = 1$ (resp. $\mathfrak{b}_q(u_i, v_i) = 1$) for every $i$; and (iii) all other pairs of vectors in $\mathcal{B}$ are orthogonal. The 2-dimensional hyperbolic bilinear form is called the *hyperbolic plane* and is denoted by $\mathbb{H}_\lambda$.

The group of all isometries of a quadratic space $(V, q)$, i.e., the *orthogonal group* of $(V, q)$ is denoted by $O(V, q)$. An isometry $\tau \in O(V, q)$ is called an *involution* if $\tau^2 = \mathrm{id}$. The *spinor norm* of an isometry $\tau$ is denoted by $\theta(\tau)$ (see [4, p. 76 and p. 137]). The *fixed space* and the *residual space* of an isometry $\tau$ are defined as follows:

$$\operatorname{Fix}(\tau) = \{x \in V \mid \tau(x) = x\} \quad \text{and} \quad \operatorname{Res}(\tau) = \{\tau(x) - x \mid x \in V\}.$$

Note that $\dim_F \operatorname{Res}(\tau) + \dim_F \operatorname{Fix}(\tau) = \dim_F V$. Also, it is readily seen that $\operatorname{Res}(\tau) = \operatorname{Fix}(\tau)^\perp$. The map $\omega_\tau : \operatorname{Res}(\tau) \times \operatorname{Res}(\tau) \to F$ defined by

$$\omega_\tau(\tau(x) - x, \tau(y) - y) = \mathfrak{b}_q(\tau(x) - x, y), \quad \text{for } x, y \in V,$$

is called the *Wall form* of $\tau$. This map is a nondegenerate bilinear form satisfying $\omega_\tau(u,u) = -q(u)$ for $u \in \mathrm{Res}(\tau)$ (see [5, (1.1)]). Also, as stated in [5, p. 116], this form is symmetric (resp. antisymmetric) if and only if $\tau^2 = \mathrm{id}$ (resp. $(\tau - \mathrm{id})^2 = 0$).

For $u \in V$ with $q(u) \neq 0$, the involution $\tau_u \in O(V,q)$ defined by $\tau_u(x) = x - \frac{\mathfrak{b}_q(u,x)}{q(u)}u$ is called the (*orthogonal*) *reflection* along $u$. Also, if $x \in V$ is an isotropic vector and $w \in (Fx)^\perp$, the *Eichler transformation* $E_{x,w} : V \to V$ is defined by

$$E_{x,w}(v) = v + \mathfrak{b}_q(v,x)w - \mathfrak{b}_q(v,w)x - q(w)\mathfrak{b}_q(v,x)x.$$

The map $E_{x,w}$ is a unipotent isometry of $(V,q)$ with $\mathrm{Res}(E_{x,w}) = Fx + Fw$.

Let $A$ be a central simple algebra over a field $F$. An *involution* on $A$ is an antiautomorphism of $A$ of order 2. We say that $\sigma$ is of the *first kind* if $\sigma|_F = \mathrm{id}$. Involutions of the first kind are either *symplectic* or *orthogonal* (see [9, (2.5)]). The *discriminant* of an orthogonal involution $\sigma$ is denoted by $\mathrm{disc}\,\sigma$ (see [9, (7.2)]). Also, the set of *alternating* and *symmetric* elements of $(A,\sigma)$ are defined as follows:

$$\mathrm{Alt}(A,\sigma) = \{a - \sigma(a) \mid a \in A\} \quad \text{and} \quad \mathrm{Sym}(A,\sigma) = \{a \in A \mid \sigma(a) = a\}.$$

## 3. Decomposition in terms of the Wall form

**Lemma 3.1.** *For an isometry $\tau$ of a quadratic space $(V,q)$ the following conditions are equivalent:* (1) $(\tau - \mathrm{id})^2 = 0$. (2) $\mathrm{Res}(\tau) \subseteq \mathrm{Fix}(\tau)$. (3) *The restriction of $q$ to* $\mathrm{Res}(\tau)$ *is totally singular.*

*Proof.* See [4, pp. 46-47] (note that the argument given there works in arbitrary characteristic). □

**Lemma 3.2.** *Let $(V,q)$ be a quadratic space over a field $F$ and let $\tau$ be an isometry of $(V,q)$ with $(\tau - \mathrm{id})^2 = 0$. If $W$ is a complement of* $\mathrm{Res}(\tau)$ *in* $\mathrm{Fix}(\tau)$*, then*

(1) *$W$ is regular and $\tau = \mathrm{id}_W \perp \tau|_{W^\perp}$.*
(2) $\dim_F W^\perp = 2\dim_F \mathrm{Res}(\tau)$.
(3) $\mathrm{Fix}(\tau|_{W^\perp}) = \mathrm{Res}(\tau|_{W^\perp}) = \mathrm{Res}(\tau)$.

*Proof.* (1) Let $u \in W \cap W^\perp$ and let $x$ be an arbitrary vector in $\mathrm{Fix}(\tau) = W \oplus \mathrm{Res}(\tau)$. Write $x = w + v$ for some $w \in W$ and $v \in \mathrm{Res}(\tau)$. Since $u \in W^\perp$, we have $\mathfrak{b}_q(u,w) = 0$. Also, as $u \in W \subseteq \mathrm{Fix}(\tau)$ and $v \in \mathrm{Res}(\tau) = \mathrm{Fix}(\tau)^\perp$, we get $\mathfrak{b}_q(u,v) = 0$. Thus, $\mathfrak{b}_q(u,x) = 0$, i.e., $u \in \mathrm{Fix}(\tau)^\perp = \mathrm{Res}(\tau)$. The equality $W \cap \mathrm{Res}(\tau) = \{0\}$ then implies that $u = 0$, i.e., $W$ is regular. By [3, (7.22)] we have $V = W \perp W^\perp$, hence $\tau = \tau|_W \perp \tau|_{W^\perp} = \mathrm{id}_W \perp \tau|_{W^\perp}$.
(2) Since $\dim_F W^\perp = \dim_F V - \dim_F W$, we get

$$\dim_F W^\perp = \dim_F V - (\dim_F \mathrm{Fix}(\tau) - \dim_F \mathrm{Res}(\tau)) = 2\dim_F \mathrm{Res}(\tau).$$

(3) Set $\tau' = \tau|_{W^\perp}$. Then $\mathrm{Res}(\tau) = \mathrm{Res}(\mathrm{id}_W) \perp \mathrm{Res}(\tau') = \mathrm{Res}(\tau')$. Using the second part, the relation $\dim_F \mathrm{Res}(\tau') + \dim_F \mathrm{Fix}(\tau') = \dim_F W^\perp$ yields $\dim_F \mathrm{Fix}(\tau') = \dim_F \mathrm{Res}(\tau')$. This, together with (3.1) implies that $\mathrm{Fix}(\tau') = \mathrm{Res}(\tau')$. □

**Definition 3.3.** ([18]) Let $(V,q)$ be a 4-dimensional quadratic space over a field $F$. An isometry $\tau \in O(V,q)$ is called an *interchange isometry* if $\mathrm{Fix}(\tau)$ is a 2-dimensional totally isotropic subspace of $(V,q)$.

**Remarks 3.4.** (1) If $\tau$ is an interchange isometry, then $(V,q)$ is hyperbolic by [3, (7.28)]. Also, dimension count implies that $\mathrm{Fix}(\tau) = \mathrm{Fix}(\tau)^\perp = \mathrm{Res}(\tau)$. In particular, $(\tau - \mathrm{id})^2 = 0$ by (3.1).
(2) If $\mathrm{char}\, F \neq 2$, then (3.1) shows that for a 4-dimensional isometry $\tau$ the following conditions are equivalent: (i) $\tau$ is an interchange isometry. (ii) $(\tau - \mathrm{id})^2 = 0$ and $\tau \neq \mathrm{id}$. (iii) $\mathrm{Fix}(\tau)^\perp = \mathrm{Fix}(\tau)$.

**Proposition 3.5.** *Let $(V,q)$ be a 4-dimensional quadratic space over a field $F$. For an isometry $\tau$ of $(V,q)$ the following conditions are equivalent:*

(1) *$\tau$ is an interchange isometry.*
(2) *There exists a hyperbolic basis $(x,y,w,z)$ of $(V,q)$ such that $(x,w)$ is a basis of $\mathrm{Fix}(\tau)$, $\tau(y)=y+w$ and $\tau(z)=z-x$.*
(3) *$(\tau-\mathrm{id})^2=0$ and $\tau$ is indecomposable.*

*Proof.* (1) $\Rightarrow$ (2) By [13, (3.5)], it suffices to consider the case where $\mathrm{char}\, F \neq 2$. Since $\mathrm{Res}(\tau)$ is singular and $\dim_F \mathrm{Res}(\tau)=2$, $\tau$ is an Eichler transformation by [10, (3.10)]. Write $\tau = E_{x,w}$ for some $x,w \in \mathrm{Res}(\tau)=\mathrm{Fix}(\tau)$ and extend $(x,w)$ to a hyperbolic basis $(x,y,w,z)$ of $(V,q)$. Then $\tau(y)=E_{x,w}(y)=y+w$ and $\tau(z)=E_{x,w}(z)=z-x$, hence $(x,y,w,z)$ is the desired basis.

(2) $\Rightarrow$ (3) Since $\mathrm{Res}(\tau)=Fx+Fw$ is totally isotropic, the equality $(\tau-\mathrm{id})^2=0$ follows from (3.1). If $\mathrm{char}\, F \neq 2$, then [15, (247.1)] shows that $\tau$ is indecomposable. Suppose that $\mathrm{char}\, F=2$. Then every proper regular subspace $V_1$ of $V$ is 2-dimensional over $F$. If in addition $\tau(V_1)=V_1$, then $\tau|_{V_1}$ is either the identity on $V_1$ or a reflection (see [18, Lem. 1]). In both cases $\mathrm{Fix}(\tau|_{V_1})$ is regular, contradicting the fact that $\mathrm{Fix}(\tau|_{V_1}) \subseteq \mathrm{Fix}(\tau)=Fx+Fw$ is totally isotropic.

(3) $\Rightarrow$ (1) Observe first that $\mathrm{Res}(\tau)=\mathrm{Fix}(\tau)$, since otherwise by (3.2) every complement of $\mathrm{Res}(\tau)$ in $\mathrm{Fix}(\tau)$ would be regular and $\tau$-invariant, contradicting the hypothesis. Since $\dim_F \mathrm{Res}(\tau)+\dim_F \mathrm{Fix}(\tau)=\dim_F V=4$, we obtain $\dim_F \mathrm{Fix}(\tau)=2$. If $\mathrm{char}\, F \neq 2$, then $\mathrm{Fix}(\tau)$ is totally isotropic by (3.1), hence $\tau$ is an interchange isometry. Suppose that $\mathrm{char}\, F=2$ and let $u$ be an arbitrary vector in $\mathrm{Fix}(\tau)=\mathrm{Res}(\tau)$. We claim that $q(u)=0$. Choose $v \in V$ such that $u=\tau(v)-v$. Then $Fu+Fv$ is a $\tau$-invariant subspace of $V$. If $q(u)\neq 0$, then $\mathfrak{b}_q(u,v)=\omega_\tau(u,u)=-q(u)\neq 0$, i.e., $Fu+Fv$ is regular, a contradiction. Thus, $q(u)=0$ and $\tau$ is an interchange isometry. □

**Remark 3.6.** Let $(V,q)$ be a 4-dimensional quadratic space over a field $F$ and let $\tau \in O(V,q)$ be an interchange isometry. If $(x,y,w,z)$ is a basis of $V$ with the properties stated in (3.5), then a direct computation shows that $\tau$ coincides with the restriction to $V$ of the inner automorphism of $C(q)$ induced by $g:=1+wx \in C(q)$. Furthermore, if $\mathrm{char}\, F=2$, the element $g$ can bo chosen more naturally as follows: by [13, (3.6)], there exist 2-dimensional regular subspaces $V_1$ and $V_2$ of $V$ such that $V=V_1 \perp V_2$, $\tau(V_1)=V_2$ and $\tau(V_2)=V_1$. Let $(u_1,v_1)$ be any basis of $V_1$ with $\mathfrak{b}_q(u_1,v_1)=1$. Then the inner automorphism of $C(q)$ induced by

$$g=1+(u_1+\tau(u_1))(v_1+\tau(v_1)) \in C(q),$$

restricts to $\tau$ on $V$. The element $g$ corresponds to the *Goldman element* of $C(q|_{V_1})\otimes C(q|_{V_1})$ (see [9, p. 35 and Exercise 8, p. 63]).

**Remarks 3.7.** Let $(V,q)$ be a quadratic space over a field $F$ and let $\tau$ be an isometry of $(V,q)$ with $(\tau-\mathrm{id})^2=0$. Consider the Wall form $\omega_\tau$.
(1) If $\mathrm{char}\, F \neq 2$, then $\omega_\tau$ is an antisymmetric bilinear form, hence it is alternating. By [3, (1.8)], $\omega_\tau$ is isometric to $n\mathbb{H}_{-1}$ for some $n$.
(2) If $\mathrm{char}\, F=2$, then $\tau$ is an involution in $O(V,q)$. Thus, $\omega_\tau$ is symmetric, which is either nonalternating or isometric to $n\mathbb{H}_{-1}$ for some $n$ (see [3, (1.8)]).

**Lemma 3.8.** *Let $(V,q)$ be a quadratic space over a field $F$ of characteristic 2 and let $\tau \in O(V,q)$. For a vector $u \in V$ the following conditions are equivalent:*

(1) *There exists a regular subplane $V'$ of $V$ containing $u$ such that $\tau|_{V'}=\tau_u$.*
(2) *$u \in \mathrm{Res}(\tau)$ and $\omega_\tau(u,u)\neq 0$.*

*Proof.* (1) $\Rightarrow$ (2) Choose a vector $v \in V'$ such that $\mathfrak{b}_q(u,v)=-q(u)\in F^\times$. Then $\tau(v)-v=u$, hence $u \in \mathrm{Res}(\tau)$. We also have $\omega_\tau(u,u)=-q(u)\neq 0$.

(2) ⇒ (1) Write $u = \tau(v) - v$ for some $v \in V$ and set $V' = Fu + Fv$. Then $\mathfrak{b}_q(u, v) = \omega_\tau(u, u) = -q(u) \neq 0$, hence $V'$ is regular. Since $\tau(u) = u$ and $\tau(v) = v + u = v - \frac{\mathfrak{b}_q(u,v)}{q(u)}u$, we have $\tau|_{V'} = \tau_u$. □

**Lemma 3.9.** *Let $(V, q)$ be a quadratic space over a field $F$ and let $\tau$ be an isometry of $(V, q)$ with $(\tau - \mathrm{id})^2 = 0$. For a 2-dimensional subspace $U$ of $V$ the following conditions are equivalent:*

(1) *There exists a 4-dimensional regular subspace $V'$ of $V$ such that $\tau|_{V'}$ is an interchange isometry with $\mathrm{Res}(\tau|_{V'}) = U$.*
(2) *$U \subseteq \mathrm{Res}(\tau)$ and $\omega_\tau|_U \simeq \mathbb{H}_{-1}$.*

*Proof.* (1) ⇒ (2) Choose a hyperbolic basis $(x, y, w, z)$ of $V'$ with the properties stated in (3.5). Then $U = Fx + Fw$, $\omega_\tau(w, w) = -q(w) = 0$ and $\omega_\tau(x, x) = -q(x) = 0$. Also, the equalities $w = \tau(y) - y$ and $x = z - \tau(z)$ imply that $\omega_\tau(x, w) = \mathfrak{b}_q(x, y) = 1$ and $\omega_\tau(w, x) = \mathfrak{b}_q(w, -z) = -1$, hence $\omega_\tau|_U \simeq \mathbb{H}_{-1}$.

(2) ⇒ (1) Let $(x, w)$ be a hyperbolic basis of $U \subseteq \mathrm{Res}(\tau)$. Write $w = \tau(y) - y$ and $x = z - \tau(z)$ for some $y, z \in V$. Set $V' := Fx + Fy + Fw + Fz$. The relation $(\tau - \mathrm{id})^2 = 0$ implies that $\tau(x) = x$ and $\tau(w) = w$, hence $V'$ is a $\tau$-invariant subspace of $V$ and $\mathrm{Fix}(\tau|_{V'}) = \mathrm{Res}(\tau|_{V'}) = U$. By (3.1) we have $\mathfrak{b}_q(x, w) = 0$. Also, $q(w) = -\omega_\tau(w, w) = 0$ and $q(x) = -\omega_\tau(x, x) = 0$, so $\mathrm{Fix}(\tau|_{V'})$ is a 2-dimensional totally isotropic subspace of $V'$. It remains to show that $V'$ is regular. Let $u \in V' \cap V'^{\perp}$. Since $\tau(V') = V'$ one has $\tau(u) - u \in V' \cap V'^{\perp}$. On the other hand we have $\tau(u) - u \in \mathrm{Res}(\tau|_{V'}) = Fx + Fw$, hence there exists $a, b \in F$ such that $\tau(u) - u = ax + aw$. Then

$$a = \omega_\tau(ax + bw, w) = \omega_\tau(\tau(u) - u, w) = \mathfrak{b}_q(\tau(u) - u, y) = 0.$$

Similarly, we have $b = 0$. Thus, $\tau(u) = u$, i.e., $u \in \mathrm{Fix}(\tau|_{V'}) = U = Fx + Fw$. Repeating the above argument we obtain $u = 0$, hence $V'$ is regular. □

**Theorem 3.10.** *Let $(V, q)$ be a quadratic space over a field $F$ and let $\tau \in O(V, q)$ with $(\tau - \mathrm{id})^2 = 0$. Let $W$ be a complement of $\mathrm{Res}(\tau)$ in $\mathrm{Fix}(\tau)$. Then there exists a decomposition $V = W \perp V_1 \perp \cdots \perp V_m$ into $\tau$-invariant regular subspaces, where $\tau|_W = \mathrm{id}$ and the integer $m$ and the subspaces $V_i$ can be described as follows:*

(1) *If $\omega_\tau$ is alternating, then $m = \frac{1}{2}\dim_F \mathrm{Res}(\tau)$, $\dim_F V_i = 4$ and $\tau|_{V_i}$ is an interchange isometry for every $i$. More precisely, if $(x_1, w_1, \cdots, x_m, w_m)$ is a hyperbolic basis of $(\mathrm{Res}(\tau), \omega_\tau) \simeq m\mathbb{H}_{-1}$, then the subspace $V_i$, $i = 1, \cdots, m$, can be chosen such that $\mathrm{Res}(\tau|_{V_i}) = Fx_i + Fw_i$.*
(2) *If $\omega_\tau$ is nonalternating, then $m = \dim_F \mathrm{Res}(\tau)$, $\dim_F V_i = 2$ and $\tau|_{V_i}$ is a reflection for every $i$. More precisely, if $(u_1, \cdots, u_m)$ is an orthogonal basis of $(\mathrm{Res}(\tau), \omega_\tau)$, then the subspace $V_i$, $i = 1, \cdots, m$, can be chosen such that $\tau|_{V_i} = \tau_{u_i}$.*

*Proof.* Set $s = \dim_F \mathrm{Res}(\tau)$, $U = W^\perp$, $q' = q|_U$ and $\tau' = \tau|_U$. By (3.2), $\tau = \mathrm{id}_W \perp \tau'$, $\mathrm{Res}(\tau') = \mathrm{Res}(\tau)$ and $\dim_F U = 2s$. We also have $(\tau' - \mathrm{id})^2 = 0$ and $\omega_{\tau'} = \omega_\tau$.

(1) Set $m = \frac{s}{2}$. Then $\dim_F U = 4m$. By (3.9) there exists a 4-dimensional subspace $V_1 \subseteq U$ such that $\tau'|_{V_1}$ is an interchange isometry and $\mathrm{Res}(\tau'|_{V_1}) = Fx_1 + Fw_1$. Let $U'$ be the orthogonal complement of $V_1$ in $U$ with respect to $q'$. Then $\dim_F U' = 4(m - 1)$. For $i = 2, \cdots, m$ and $v \in V_1$, we have $\mathfrak{b}_q(x_i, v) = \omega_\tau(x_i, \tau(v) - v) = 0$, since $\tau(v) - v \in \mathrm{Res}(\tau'|_{V'}) = Fx_1 + Fw_1$. Similarly, we have $\mathfrak{b}_q(w_i, v) = 0$, hence $x_i, w_i \in U'$. It follows that $(x_2, w_2, \cdots, x_m, w_m)$ is a hyperbolic basis of $\omega_{\tau'|_{U'}} \simeq (m - 1)\mathbb{H}_{-1}$. By induction and dimension count one can write

$$U = V_1 \perp \cdots \perp V_m,$$

where every $V_i$ is 4-dimensional subspaces of $U$ and $\tau'|_{V_i}$ is an interchange isometry with $\mathrm{Res}(\tau'|_{V_1}) = Fx_1 + Fw_1$. Thus, $V = W \perp U = W \perp V_1 \perp \cdots \perp V_m$ is the desired decomposition.

(2) By (3.7 (2)) we have $\operatorname{char} F = 2$. Set $m = s$. In view of (3.8), there exists a 2-dimensional subspace $V_1 \subseteq U$ containing $U$ such that $\tau'|_{V_1} = \tau_{u_1}$. Let $U'$ be the orthogonal complement of $V_1$ in $U$. For $i = 2, \cdots, n$, we have $u_i \in U'$, since $\mathfrak{b}_q(u_i, v_1) = \omega_\tau(u_i, u_1) = 0$. Thus, $(u_2, \cdots, u_n)$ is an orthogonal basis of $\omega_{\tau'|_{U'}}$. The result again follows from induction and dimension count. □

**Remark 3.11.** If $\operatorname{char} F = 2$, (3.10) gives an alternative proof of the Wiitala decomposition of involutions in orthogonal group (see [18, Theorem. 1]). Also, the subspace $W$ in (3.10) is a Wiitala subspace of $(V, \tau)$ in the sense of [13, (3.8)]. Finally, $\tau$ is of *interchanging kind* if and only if $\omega_\tau$ is alternating (see [13, (3.19)]).

## 4. The induced involution on Clifford algebra in characteristic two

Throughout this section, $F$ denotes a field of characteristic 2.

Let $(V, q)$ be a quadratic space over $F$. Every isometry $\tau$ of $(V, q)$ with $(\tau - \mathrm{id})^2 = 0$ is an involution in the orthogonal group. Such an involution induces an involution $J_\tau$ on $C(q)$ satisfying $J_\tau(v) = \tau(v)$ for $v \in V$. The aim of this section is to study the relation between the Wall form $\omega_\tau$ and some invariants of $(C(q), J_\tau)$, introduced in [2] and [12]. We first recall some constructions from [2] and [12].

An algebra with involution $(A, \sigma)$ is called *totally decomposable* if it decomposes as tensor products of *quaternion algebras* (i.e., central simple algebras of degree 2) with involution. Let $(A, \sigma)$ be a totally decomposable algebra of degree $2^n$ with orthogonal involution over $F$. As proved in [12, pp. 10-11], there exists a unique, up to isomorphism, subalgebra $S \subseteq \mathrm{Sym}(A, \sigma)$ such that (i) $x^2 \in F$ for $x \in S$; (ii) $\dim_F S = \deg_F A$; (iii) $S$ is self-centralizing, i.e., $C_A(S) = S$; and (iv) $S$ is generated as an $F$-algebra by exactly $n$ elements. The isomorphism class of $S$ is denoted by $\Phi(A, \sigma)$. Furthermore, the algebra $\Phi(A, \sigma)$ has a *set of alternating generators*, i.e., a set $\{u_1, \cdots, u_n\} \subseteq \mathrm{Alt}(A, \sigma)$ consisting of pairwise commutative square-central units satisfying $u_{i_1} \cdots u_{i_l} \in \mathrm{Alt}(A, \sigma)$ for every $1 \leq l \leq n$ and $1 \leq i_1 < \cdots < i_l \leq n$. If $\{u_1, \cdots, u_n\}$ is a set of alternating generators of $\Phi(A, \sigma)$, then $\Phi(A, \sigma) \simeq F[u_1, \cdots, u_n]$.

Let $(A, \sigma) \simeq \bigotimes_{i=1}^n (Q_i, \sigma_i)$ be a decomposition of $A$ into quaternion algebras with involution and let $\alpha_i \in F^\times$, $i = 1, \cdots, n$, be a representative of the class $\operatorname{disc} \sigma_i \in F^\times / F^{\times 2}$. The form $\langle\langle \alpha_1, \cdots, \alpha_n \rangle\rangle$ is called the *Pfister invariant* of $(A, \sigma)$ and is denoted by $\mathfrak{Pf}(A, \sigma)$. By [2, (7.5)], $\mathfrak{Pf}(A, \sigma)$ is independent of the decomposition of $(A, \sigma)$. Finally, if $\{u_1, \cdots, u_n\}$ is a set of alternating generators of $\Phi(A, \sigma)$ with $u_i^2 = \alpha_i \in F^\times$, $i = 1, \cdots, n$, then $\mathfrak{Pf}(A, \sigma) \simeq \langle\langle \alpha_1, \cdots, \alpha_n \rangle\rangle$ (see [12, pp. 11-12]).

Our first result is the following restatement of [13, (4.7)], in view of (3.11):

**Lemma 4.1.** *Let $(V, q)$ be a quadratic space over $F$ and let $\tau \in O(V, q)$ be an involution. The involution $J_\tau$ on $C(q)$ is orthogonal if and only if* $\mathrm{Res}(\tau) = \mathrm{Fix}(\tau)$.

**Lemma 4.2.** *Let $(V, q)$ be a 2-dimensional quadratic space over $F$ and let $\tau \in O(V, q)$ be a reflection. Then $(C(q), J_\tau) \simeq (M_2(F), t)$ if and only if $\theta(\tau)$ is trivial.*

*Proof.* If $(C(q), J_\tau) \simeq (M_2(F), t)$, then $\operatorname{disc} J_\tau$ is trivial, hence $\theta(\tau) = 1$ by [13, (4.11)]. Conversely, if $\theta(\tau)$ is trivial, then $q$ represents 1. Thus, there exists $x \in V$ such that $x^2 = 1 \in C(q)$. Then $(x + 1)^2 = 0$, hence $C(q)$ splits. Also, [13, (4.11)] shows that $\operatorname{disc} \sigma = \theta(\tau)$ is trivial, hence $(C(q), J_\tau) \simeq (M_2(F), t)$ by [9, (7.4)]. □

The next result follows from [13, (6.10)] and (4.2).

**Corollary 4.3.** *Let $(V, q)$ be a 4-dimensional quadratic space over $F$ and let $\tau$ be an interchange isometry of $(V, q)$. Then $(C(q), J_\tau) \simeq (M_4(F), t)$.*

**Corollary 4.4.** *Let $(V,q)$ be a quadratic space over $F$ and let $\tau \in O(V,q)$ be an involution. Let $W$ be a complement of $\mathrm{Res}(\tau)$ in $\mathrm{Fix}(\tau)$. If $\omega_\tau$ is alternating, then $(C(q),J_\tau) \simeq (C(q|_W),J_{\mathrm{id}}) \otimes_F (M_{2^s}(F),t)$, where $s = \dim_F \mathrm{Res}(\tau)$.*

*Proof.* The result follows from (3.10 (1)), [13, (4.4)] and (4.3). □

**Lemma 4.5.** *Let $(V,q)$ be a quadratic space over $F$. If $\tau$ is an involution in $(V,q)$, then $(C(q),J_\tau)$ is totally decomposable.*

*Proof.* By (3.10), there exists a decomposition $V = W \perp V_1 \perp \cdots \perp V_m$ into regular subspaces such that $\tau|_W = \mathrm{id}$ and either $\dim_F V_i = 4$ and $\tau_i := \tau|_{V_i}$ is an interchange isometry, or $\dim_F V_i = 2$ and $\tau_i$ is a reflection. By [13, (4.4)] we have

$$(C(q),J_\tau) \simeq (C(q|_W),J_{\mathrm{id}}) \otimes (C(q|_{V_1}),J_{\tau_1}) \otimes \cdots \otimes (C(q|_{V_m}),J_{\tau_m}).$$

If $\tau_i$ is a reflection, then $(C(q|_{V_i}),J_{\tau_i})$ is a quaternion algebra with involution. Thus, in view of (4.3), it suffices to show that $(C(q|_W),J_{\mathrm{id}})$ is totally decomposable. Since $W$ is regular, one can write $W = W_1 \perp \cdots \perp W_k$, where $k = \frac{1}{2}\dim_F W$ and every $W_i$ is a 2-dimensional regular subspace of $W$. Thus, $(C(q|_W),J_{\mathrm{id}}) \simeq \bigotimes_{i=1}^k (C(q|_{W_i}),J_{\mathrm{id}})$ is totally decomposable. □

The next result states the relation between the subalgebra $\Phi(C(q),J_\tau)$ and the Wall form of $\tau$. Recall that if $(V,\mathfrak{b})$ is a symmetric bilinear space over a field $F$, the map $\varphi_\mathfrak{b} : V \to F$ defined by $\varphi_\mathfrak{b}(v) = \mathfrak{b}(v,v)$ is a quadratic form, called the *associated quadratic form* of $\mathfrak{b}$.

**Proposition 4.6.** *Let $(V,q)$ be a quadratic space over $F$ and let $\tau \in O(V,q)$ be an involution such that $\mathrm{Res}(\tau) = \mathrm{Fix}(\tau)$. If $\varphi$ is the associated quadratic form of $\omega_\tau$, then $\Phi(C(q),J_\tau) \simeq C(\varphi) \simeq C(q|_{\mathrm{Res}(\tau)})$.*

*Proof.* Since $\varphi(v) = \omega_\tau(v,v) = -q(v)$ for every $v \in \mathrm{Res}(\tau)$, we have $\varphi = -q|_{\mathrm{Res}(\tau)}$. Thus, $C(q|_{\mathrm{Res}(\tau)}) \simeq C(\varphi)$. By (3.1), $\varphi$ is totally singular, hence $C(\varphi)$ is commutative. Let $(u_1,\cdots,u_s)$ be a basis of $\mathrm{Res}(\tau)$ over $F$, where $s = \dim_F \mathrm{Res}(\tau) = \frac{1}{2}\dim_F V$. Since for every $i$, $u_i \in \mathrm{Sym}(C(q),J_\tau)$ and $u_i^2 \in F$, the commutativity of $C(\varphi)$ implies that $C(\varphi) \subseteq \mathrm{Sym}(C(q),J_\tau)$ and $x^2 \in F$ for every $x \in C(\varphi)$. Also, $\dim_F C(\varphi) = 2^s$ and $C(\varphi)$ is generated as an $F$-algebra by $u_1,\cdots,u_s$, hence $C(\varphi)$ is a Frobenius algebra by [12, (3.4)]. Finally, as $\dim_F C(\varphi) = 2^s = \deg_F C(q)$, $C(\varphi) \subseteq C(q)$ is self-centralizing by [8, (2.2.3 (1))], thus $C(\varphi) \simeq \Phi(C(q),J_\tau)$. □

**Theorem 4.7.** *Let $(V,q)$ be a quadratic space over $F$ and let $\tau \in O(V,q)$ be an involution such that $\mathrm{Res}(\tau) = \mathrm{Fix}(\tau)$. Let $s = \dim_F \mathrm{Res}(\tau) = \frac{1}{2}\dim_F V$.*

1. *If $\omega_\tau$ is alternating then $(C(q),J_\tau) \simeq (M_{2^s}(F),t)$ and $\mathfrak{Pf}(C(q),J_\tau) \simeq \langle\!\langle 1,\cdots,1\rangle\!\rangle$.*
2. *If $\omega_\tau$ is nonalternating then every orthogonal basis of $\mathrm{Res}(\tau)$ is a set of alternating generators of $\Phi(C(q),J_\tau)$. In particular, if $\omega_\tau \simeq \langle\alpha_1,\cdots,\alpha_s\rangle$ for $\alpha_1,\cdots,\alpha_s \in F^\times$ then $\mathfrak{Pf}(C(q),J_\tau) \simeq \langle\!\langle \alpha_1,\cdots,\alpha_s\rangle\!\rangle$.*

*Proof.* The first part follows from (4.4) and [12, (5.7)]. To prove the second part let $(u_1,\cdots,u_s)$ be an orthogonal basis of $\mathrm{Res}(\tau)$. Then for $i = 1,\cdots,s$, we have $u_i^2 = q(u_i) \in F$ in $C(q)$. The equality $\mathfrak{b}_q(u_i,u_j) = 0$ implies that $u_iu_j = u_ju_i \in C(q)$ for every $i,j$. By (3.10 (1)) one can write $\tau = \tau_{u_1} \perp \cdots \perp \tau_{u_s}$, where every $\tau_{u_i}$ is a reflection of a 2-dimensional $\tau$-invariant subspaces $V_i$ of $V$. For $i = 1,\cdots,s$, choose an element $v_i \in V_i$ such that $u_i = v_i - \tau(v_i)$. Then

$$u_{i_1}\cdots u_{i_l} = v_{i_1}u_{i_2}\cdots u_{i_l} - J_\tau(v_{i_1}u_{i_2}\cdots u_{i_l}) \in \mathrm{Alt}(C(q),J_\tau),$$

for every $1 \le i_1 < \cdots < i_l \le s$ and $1 \le l \le s$. Since $\mathrm{Res}(\tau) = \mathrm{Fix}(\tau)$, we have $\dim_F V = 2s$, hence $\deg_F C(q) = 2^s$ and $\{u_1,\cdots,u_s\}$ is a set of alternating generators of $\Phi(C(q),J_\tau)$. □

The proof of the following result is left to the reader.

**Lemma 4.8.** *If $X$ is a symmetric matrix with $X^2 \in F$, then $X^2 \in F^2$.*

**Theorem 4.9.** *Let $(V, q)$ be a $2n$-dimensional quadratic space over $F$. For an involution $\tau$ of $(V, q)$ with $\mathrm{Res}(\tau) = \mathrm{Fix}(\tau)$ the following conditions are equivalent:*

(1) $(C(q), J_\tau) \simeq (M_{2^n}(F), t)$.
(2) $q(x) \in F^2$ *for every* $x \in \mathrm{Res}(\tau)$.
(3) $\omega_\tau(x, x) \in F^2$ *for every* $x \in \mathrm{Res}(\tau)$.
(4) *Either* $\omega_\tau \simeq n\mathbb{H}_{-1}$ *or* $\omega_\tau \simeq n\langle 1\rangle$.

*Proof.* Let $f : (C(q), J_\tau) \simeq (M_{2^n}(F), t)$ be an isomorphism of algebras with involution. For every $x \in \mathrm{Res}(\tau)$, $f(x)$ is a symmetric matrix, since $\tau(x) = x$. As $f(x)^2 = q(x) \in F$, we get $f(x)^2 \in F^2$ by (4.8), i.e., $q(x) \in F^2$. This proves (1) $\Rightarrow$ (2).

The implications (2) $\Rightarrow$ (3) and (3) $\Rightarrow$ (4) are evident. To prove (4) $\Rightarrow$ (1) observe that in view of (4.4), it suffices to consider the case where $\omega_\tau \simeq n\langle 1\rangle$. Let $(u_1, \cdots, u_n)$ be an orthogonal basis of $\omega_\tau$. By (3.10 (2)) one can write $V = V_1 \perp \cdots \perp V_n$, where every $V_i$ is a 2-dimensional subspace of $V$ containing $u_i$ and $\tau|_{V_i} = \tau_{u_i}$. Thus, [13, (4.4)] and (4.2) imply that

$$(C(q), J_\tau) \simeq \bigotimes_{i=1}^{n} (C(q|_{V_i}), J_{\tau_{u_i}}) \simeq \bigotimes_{i=1}^{n} (M_2(F), t) \simeq (M_{2^n}(F), t). \qquad \square$$

## References


[1] E. A. Connors, The structure of $\mathrm{O}'(V)/\mathrm{DO}(V)$ in the defective case. *J. Algebra* **34** (1975), 74–83.

[2] A. Dolphin, Orthogonal Pfister involutions in characteristic two. *J. Pure Appl. Algebra* **218** (2014), no. 10, 1900–1915.

[3] R. Elman, N. Karpenko, A. Merkurjev, *The algebraic and geometric theory of quadratic forms.* American Mathematical Society Colloquium Publications, 56. American Mathematical Society, Providence, RI, 2008.

[4] L. C. Grove, *Classical Groups and Geometric Algebra*, Graduate Studies in Mathematics, 39. American Mathematical Society, Providence, RI, 2002.

[5] A. Hahn, Unipotent elements and the spinor norms of Wall and Zassenhaus. *Arch. Math.* **32** (1979), no. 2, 114–122.

[6] A. Hahn, The elements of the orthogonal group $\Omega_n(V)$ as products of commutators of symmetries. *J. Algebra* **184** (1996), no. 3, 927–944.

[7] A. Hahn, The Zassenhaus decomposition for the orthogonal group: properties and applications. Proceedings of the Conference on Quadratic Forms and Related Topics (Baton Rouge, LA, 2001). *Doc. Math.* 2001, Extra Vol., 165–181.

[8] N. Jacobson, *Finite-dimensional division algebras over fields.* Springer-Verlag, Berlin, 1996.

[9] M.-A. Knus, A. S. Merkurjev, M. Rost, J.-P. Tignol, *The book of involutions.* American Mathematical Society Colloquium Publications, 44. American Mathematical Society, Providence, RI, 1998.

[10] F. Knüppel, The length-problem for Eichler-transformations. *Forum Math.* **10** (1998), no. 1, 59–74.

[11] M. G. Mahmoudi, Orthogonal symmetries and Clifford algebras. *Proc. Indian Acad. Sci. Math. Sci.* **120** (2010), no. 5, 535–561.

[12] M. G. Mahmoudi, A.-H. Nokhodkar, On totally decomposable algebras with involution in characteristic two. LAG preprint server, http://www.math.uni-bielefeld.de/LAG/man/549.html.

[13] M. G. Mahmoudi, A.-H. Nokhodkar, Involutions of a Clifford algebra induced by involutions of orthogonal group in characteristic 2. *Comm. Algebra* **43** (2015), no. 9, 3898–3919.

[14] D. B. Shapiro, *Compositions of quadratic forms.* de Gruyter Expositions in Mathematics, 33. Walter de Gruyter & Co., Berlin, 2000.

[15] E. Snapper, R. J. Troyer, *Metric affine geometry.* Second edition. Dover Books on Advanced Mathematics. Dover Publications, Inc., New York, 1989.

[16] G. E. Wall, On the conjugacy classes in the unitary, symplectic and orthogonal groups. *J. Austral. Math. Soc.* **3** (1963) 1–62.

[17] G. E. Wall, The structure of a unitary factor group. *Inst. Hautes tudes Sci. Publ. Math.* No. 1 1959 23 pp. (1959).

[18] S. A. Wiitala, Factorization of involutions in characteristic two orthogonal groups: an application of the Jordan form to group theory, *Linear Algebra Appl.*, **21** (1978), no. 1, 59–64.

[19] H. Zassenhaus, On the spinor norm. *Arch. Math.* **13** (1962) 434–451.

A.-H. Nokhodkar, a.nokhodkar@kashanu.ac.ir
Department of Pure Mathematics, Faculty of Science, University of Kashan, P. O. Box 87317-53153, Kashan, Iran.